

## The Piéla-Massako Permit/Burkina Faso - Final Report

by Wolfgang Hampel (1997), 68 pages, 55 figures and maps, 13 tables and 6 annexes

**Client:** Emerging Africa Gold Corp., Montreal/Canada      **Position held:** General Manager BF

**Scope of project:** The initial aim was to delineate an oxidized, open-pittable gold resource of at least 1 Mio ounces with a minimum GT-value of 100. The minimum target was later augmented to 2 Mio ounces.

### Work carried out:

- Installation of a permanent field camp
- compilation of all available data
- regional mapping , 1:50,000, 780 sqkm (Landsat interpretation.)
- prospect scale mapping, 1:5,000, 15 sqkm
- soil/lag/termite mound geochemistry (Au+33), 4,000 samples
- outcrop sampling, 400 samples
- ground geophysics, 25 line kilometers (VLF, MAG, IP)
- 580 m trenching, with detailed logging and sampling
- RC drilling, 4,000 m, 80 holes
- evaluation of all known orpailleur workings
- statistics of soil geochemistry
- detailed struct. and geol. interpretation of the permit
- description of known mineralisation
- development of metallogenetic models
- description and classification of targets
- description of methodology
- proposition of detailed 1-year work program
- proposition of a 1-year US \$ 1,050,000 budget


### RESULTS:

**Geological setting:** The 780 sqkm are almost entirely underlain by volcanic and volcanosedimentary sequences of Birimian age. The compositions of the metavolcanics range from intermediate to felsic, mafic to ultramafic lithologies are less abundant. The Birimian rocks are embedded in Archean gneisses and migmatites. Eburnean metagranites and metagranodiorites are frequent in the N, S and W. Mesozoic dolerite dykes crosscut the area with N115. A cluster of airborne detected magnetic anomalies in the SW are interpreted as possible kimberlites. At least three different stages of extensive duricrust formation are known.


The Birimian sequences are intensely folded and faulted. Metamorphism reaches lower greenschist facies. A major shear zone crosses the permit along its longer axis, shorter, secondary shear zones are +- parallel. Two major deformation zones, the Basieri and Madyabari 'corridors' were identified during the field campaign. They host the majority of all gold showings.

**Mineralisation:** In total 21 targets for gold, base metals and diamonds were identified. Most of the gold showings belong to the narrow vein type, though some are very high grade (up to 250 g/t Au over 0.50 m) and show considerable host rock mineralisation. Reconnaissance drilling on a 2,800 m x 250 m gold in lag anomaly returned very encouraging results with 11 m @ 5.99 g/t, 10 m @ 2,39 g/t and 22 m @ 1.24 g/t. This anomaly lacks any outcrops. Trenching points to a high-grade stratabound mineralisation in metacherts and andesitic tuffs. Many anomalies stayed 'untouched', as for example a strong Pb-Zn-Ba-anomaly that was discovered during a regional lag survey.

**Potential:** There is a fair number of prospects that could yield in excess of 200,000 oz gold each. Since 1997, the permit holder Mr. Dô Konaté has been conducting successful small-scale mining operations (gravity treatment of high-grade orpailleur rejects).



**Compilation map** showing the principal geological units, tectonic features and the main gold and diamond prospects



© 2001-2002 Wolfgang Hampel